

Various series expansions for the bilayer $S = \frac{1}{2}$ Heisenberg antiferromagnet

Zheng Weihong*

School of Physics, The University of New South Wales, Sydney, NSW 2052, Australia.

Abstract

Various series expansions have been developed for the two-layer, $S = \frac{1}{2}$, square lattice Heisenberg antiferromagnet. High temperature expansions are used to calculate the temperature dependence of the susceptibility and specific heat. At $T = 0$, Ising expansions are used to study the properties of the Néel-ordered phase, while dimer expansions are used to calculate the ground-state properties and excitation spectra of the magnetically disordered phase. The antiferromagnetic order-disorder transition point is determined to be $(J_2/J_1)_c = 2.537(5)$. Quantities computed include the staggered magnetization, the susceptibility, the triplet spin-wave excitation spectra, the spin-wave velocity, and the spin-wave stiffness. We also estimate that the ratio of the intra- and inter-layer exchange constants to be $J_2/J_1 \simeq 0.07$ for cuprate superconductor $\text{YBa}_2\text{Cu}_3\text{O}_{6.2}$.

PACS Indices: 75.10.-b., 75.10J., 75.40.Gb

I. INTRODUCTION

The magnetic properties of low dimensional systems have been the subject of intense theoretical and experimental research in recent years. It is by now well established that one-dimensional Heisenberg antiferromagnets with integer spin have a gap in the excitation spectrum, whereas those with half-integer spin have gapless excitations. The former have a finite correlation length, while for the latter it is infinite with the spin-spin correlation function decaying to zero as a power law. In 2-dimensions, the unfrustrated square-lattice Heisenberg model has long range Néel order in the ground state. It has gapless Goldstone modes as expected. In recent years much interest has focussed on systems with intermediate dimensionality and on questions of crossovers between $d = 1$ and $d = 2$, and the investigations showed that the crossover from the single Heisenberg chain to the two-dimensional antiferromagnet square lattice, obtained by assembling chains to form a “ladder” of increasing width, is far from smooth. Heisenberg ladders with an even number of legs (chain) show a completely different behavior than odd-leg-ladders. While even-leg-ladders have a spin gap and short range correlation, odd-leg-ladders have no gap and power-law correlations.

This paper is concerned with a system between 2 and 3 dimensions: the $S = \frac{1}{2}$ two-layer Heisenberg antiferromagnet, where each layer is composed of a nearest-neighbor Heisenberg model on a square lattice, and there is a further antiferromagnetic coupling between corresponding sites of each layer. The system can be described by the following Hamiltonian:

$$H = J_1 \sum_{\alpha=1,2} \sum_{\langle i,j \rangle} \mathbf{S}_{\alpha,i} \cdot \mathbf{S}_{\alpha,j} + J_2 \sum_i \mathbf{S}_{1,i} \cdot \mathbf{S}_{2,i} \quad (1)$$

where $\mathbf{S}_{\alpha,i}$ is a $S=1/2$ spin operator at site i of the α th layer, and $\langle i,j \rangle$ denotes a pair of nearest neighbor sites on a square lattice. Here J_1 is the interaction between nearest neighbor spins in one plane, and J_2 is the interaction between adjacent spins from different layers. We denote the ratio of couplings as y , that is, $y = J_2/J_1$. In the present paper, we study only the case of antiferromagnetic coupling, that is, both J_1 and J_2 are positive, although some of the series are applicable to ferromagnetic coupling case. In the small J_2/J_1 limit, the model describes two weakly interacting 2D Heisenberg antiferromagnets, where each of them is Néel ordered and possesses gapless Goldstone excitations. In the large J_2/J_1 limit, pairs of adjacent spins from different layers form spin singlets separated from triplet states by a gap, $\sim J_2$, and the system has no long-range order. Thus one expects there is a transition from ordered antiferromagnet to a quantum disordered spin liquid at a certain critical ratio of J_2/J_1 . The bilayer Heisenberg antiferromagnets has recently attracted attention for several reasons. On the experimental side, the model is relevant to the understanding of the magnetic properties of cuprate superconductors $\text{YBa}_2\text{Cu}_3\text{O}_{6+x}$ containing two weakly coupled CuO_2 layers [1–3]. On the theoretical side, it is probably the simplest unfrustrated spin system which displays a quantum disordering transition of $O(3)$ universality class, and Quantum Monte Carlo (QMC) simulation on this model do not suffer from a sign problem, making it possible to obtain precise data on the critical properties of the quantum transition. Hida [4] has performed a dimer expansion up to 6th order in J_1/J_2 , and he found the critical ratio to be $(J_2/J_1)_c = 2.56$, Gelfand [5] has recently extended the dimer expansion to 8th order, and obtains the critical ratio $(J_2/J_1)_c = 2.54(2)$. Sandvik *et al.* [6–8] have carried out a detailed QMC simulations for the finite-temperature

properties of small (J_2/J_1) and near-critical $(J_2/J_1)_c$ systems, and found a similar critical ratio: $(J_2/J_1)_c = 2.51(2)$. On the other hand, a Schwinger boson mean-field calculation by Millis and Monien [9] resulted in a very large critical ratio: $(J_2/J_1)_c = 4.48$; a self-consistent spin-wave theory [10] also predicts a large critical ratio: $(J_2/J_1)_c = 4.3$. Chubukov and Morr [10] have argued that the discrepancies between the spin-wave results and the numerical simulations (or series expansions) have a physical origin and are related to the fact that in the spin-wave approach one neglects longitudinal spin fluctuations, and after taking longitudinal spin fluctuations into account, they got the critical ratio: $(J_2/J_1)_c = 2.73$, which is reasonably close to the numerical results. The critical ratio obtained by the Schwinger-boson Gutzwiller-projection method [11] is $(J_2/J_1)_c = 3.51$.

We have carried out extensive series studies of this system, via Ising expansions and dimer expansions at $T = 0$, and also by high temperature series expansions. Our results confirm the existence of a order-disorder transition at critical coupling ratio $J_2/J_1 = 2.537(5)$. The complete spin-wave excitation spectra, spin-wave velocity, and some order parameters are computed. We also compare our results with previous calculations.

This paper is organized as follows: In Sec. II the dimer expansions are presented, and the antiferromagnetic order-disorder transition points is determined precisely. In Sec. III, the Heisenberg-Ising model is studied via the Ising expansions, which allow for estimates of the domain stability of Néel-order phase, and also allow a direct comparison with experimental data of $\text{YBa}_2\text{Cu}_3\text{O}_{6+x}$. In Sec. IV we present the results of the high-temperature expansions, and our conclusion is presented in the last section.

II. DIMER EXPANSIONS

In the limit that the exchange coupling along the rungs J_2 is much larger than the coupling J_1 within the plane, that is $y \gg 1$, the rungs interact only weakly with each other, and the dominant configuration in the ground state is the product state with the spin on each rung forming a spin singlet. The Hamiltonian in Eq. (1) can be rewritten as,

$$H/J_2 = H_0 + (1/y)V \quad (2)$$

where

$$\begin{aligned} H_0 &= \sum_i \mathbf{S}_{1,i} \cdot \mathbf{S}_{2,i} \\ V &= \sum_{\alpha=1,2} \sum_{\langle i,j \rangle} \mathbf{S}_{\alpha,i} \cdot \mathbf{S}_{\alpha,j} \end{aligned} \quad (3)$$

We can construct expansion in $1/y$ by treating the operator H_0 as the unperturbed Hamiltonian and the operator V is treated as a perturbation. The linked-cluster expansion method has been previously reviewed in several articles [12–14], and will not be repeated here.

We have carried out the dimer expansions for the $T = 0$ ground state energy per site E_0/N , the antiferromagnetic susceptibility χ , and the lowest lying triplet excitation spectrum $\Delta(k_x, k_y)$ (odd parity under interchange of the planes) up to order $(1/y)^{11}$. The resulting power series in $1/y$ for the ground state energy per site E_0/N and the antiferromagnetic

susceptibility χ are presented in Table I. Table of series coefficients for the triplet excitation spectrum $\Delta(k_x, k_y)$ would require inordinate amount of space to reproduce in print, it is available from the authors on request. Instead, we also present in Table I the series for the minimum energy gap $m = \Delta(\pi, \pi)$. The dimer expansion was firstly carried out by Hida [4] up to order $(1/y)^6$ in 1992, and extended to order $(1/y)^8$ by Gelfand [5] recently. Our results agree with these previous results, and extend the series by three terms. A list of 19093 linked clusters of up to 12 sites contribute to the triplet excitation spectrum.

To determine the critical point y_c , we construct the Dlog Padé approximants [15] to the χ and m series, and since the transition should lie in the universality class of the classical $d = 3$ Heisenberg model (our analysis also supports this), we expect that the critical index for χ and m should be approximately 1.40 and 0.71, respectively. The exponent-biased Dlog Padé approximants [5] gives the critical point $1/y_c = 0.3942(8)$, or $y_c = 2.537(5)$.

The spectra for some particular values of J_1/J_2 are illustrated in Fig. 1, where the direct sum to the series at $y = y_c$ is indeed consistent with the Padé approximants that one can construct. To compute the critical spin-wave velocity v , one expands the spectrum Δ in the vicinity of wave-vector (π, π) up to $\mathbf{k}^2$:

$$\Delta(\pi - k_x, \pi - k_y)[1/y] = C(1/y) + D(1/y)(k_x^2 + k_y^2) + \dots \quad (4)$$

and it is easy to prove [5] that the critical spin-wave velocity is equal to $(2CD)^{1/2}$ at y_c . The series coefficients for $2CD$ in J_1/J_2 are listed in Table I. Using Padé approximants, one can estimate the properties at critical point y_c : the spin-wave velocity is estimated to be $(v/J_2)^2 = 0.559(2)$, or $v/J_2 = 0.748(2)$, the ground-state energy per site is $E_0/N = -0.4456(4)J_2$, the optical spin-wave gap (the gap at $\Gamma = (0, 0)$) is $\Delta_{\text{opt}} = 1.737(1)J_2$, and the gap at $X = (\pi, 0)$ is $\Delta_X = 1.368(8)J_2$, where the uncertainty is largely associated with the uncertainty in y_c .

III. ISING EXPANSIONS

To construct an expansion about the Ising limit for this system, one has to introduce an anisotropy parameter x , and writes the Hamiltonian for Heisenberg-Ising model as:

$$H/J_1 = H_0 + xV \quad (5)$$

where

$$\begin{aligned} H_0 &= \sum_{\alpha=1,2} \sum_{\langle i,j \rangle} S_{\alpha,i}^z S_{\alpha,j}^z + y \sum_i S_{1,i}^z S_{2,i}^z + t \sum_{\alpha,i} \epsilon_{\alpha,i} S_{\alpha,i}^z \\ V &= \sum_{\alpha=1,2} \sum_{\langle i,j \rangle} (S_{\alpha,i}^x S_{\alpha,j}^x + S_{\alpha,i}^y S_{\alpha,j}^y) + y \sum_i (S_{1,i}^x S_{2,i}^x + S_{1,i}^y S_{2,i}^y) - t \sum_{\alpha,i} \epsilon_{\alpha,i} S_{\alpha,i}^z \end{aligned} \quad (6)$$

and $\epsilon_{\alpha,i} = \pm 1$ on the two sublattices. The last term in both H_0 and V is a local staggered field term, which can be included to improve convergence. The limits $x = 0$ and $x = 1$ correspond to the Ising model and the isotropic Heisenberg model respectively. The operator H_0 is taken as the unperturbed Hamiltonian, with the unperturbed ground state being the usual Néel state. The operator V is treated as a perturbation. It flips a pair of spins on neighbouring sites.

The Ising series have been calculated for the ground state energy per site E_0/N , the staggered magnetization M , the uniform perpendicular susceptibility $\chi_\perp$, and the lowest lying triplet excitation spectrum $\Delta(k_x, k_y, k_z)$ for several ratio of couplings and (simultaneously) for several values of t up to order x^{10} (the series for uniform perpendicular susceptibility $\chi_\perp$ is one order less). The resulting series for $y = 0.5, 1, 1.5, 2.5$ and $t = 0$ are listed in Tables II and III (the series coefficients for triplet excitation spectrum are listed only for case of $y = 1$), the series for other value of y are available on request.

We can also compute the spin-wave velocity v through a similar expansion of $\Delta(k_x, k_y, 0)$ in the vicinity of $k_x = k_y = 0$ as in Eq.(4), and the series coefficients for $2CD$ in x are also listed in Table II. With this series and the series for the perpendicular susceptibility $\chi_\perp$, one can get the series for the spin-wave stiffness ρ_s through the following ‘‘hydrodynamic’’ relation:

$$\rho_s = v^2 \chi_\perp . \quad (7)$$

There are two branches of the spin-wave dispersion. From the series one can see that the symmetric (optical) excitation spectrum $\Delta(k_x, k_y, 0)$ (that is, with even parity under interchange of the planes) is related to the antisymmetric (acoustic) excitation spectrum $\Delta(k_x, k_y, \pi)$ (that is, with odd parity under interchange of the planes) by

$$\Delta(k_x, k_y, 0) = \Delta(\pi - k_y, \pi - k_x, \pi) \quad (8)$$

so we only consider the antisymmetric excitation spectrum here.

The conventional linear spin-wave theory [1,2,16] predicts the magnon dispersions as:

$$\Delta(k_x, k_y, \pi) = 4SJ_1[(1 - \gamma_k^2) + y(1 + \gamma_k)/2]^{1/2} \quad (9)$$

where $\gamma_k = \frac{1}{2}[\cos(k_x) + \cos(k_y)]$. Therefore one can see that the acoustic spin-wave gap $\Delta_{\text{aco}} = \Delta(\pi, \pi, \pi)$ is zero even for nonzero J_2 , the optical gap is

$$\Delta_{\text{opt}} = \Delta(0, 0, \pi) = 4S\sqrt{J_1 J_2} \quad (10)$$

which is non-zero as long as $J_2 \neq 0$, and the gap at $X = (\pi, 0, \pi)$ is

$$\Delta_X = 4SJ_1(1 + y/2)^{1/2}. \quad (11)$$

In a recent paper, Millis and Monien [3] got the following relation for the optical spin-wave gap by using the Schwinger boson method:

$$J_2 = \frac{\Delta_{\text{opt}}^2}{16\rho_s + 2\Delta_{\text{opt}}/\pi} \quad (12)$$

and so at the limit of $J_2 \rightarrow 0$, one results that $\Delta_{\text{opt}} = 4S\sqrt{J_1 J_2 Z_\rho}$, which differs from Eq.(10) by a factor of $\sqrt{Z_\rho}$, where $Z_\rho = \rho_s/(J_1 S^2) \simeq 0.72$.

Having obtained the series we first attempt to identify critical points and to determine the nature of the phase diagram in the $(x - y)$ plane. Naive Dlog Padé analysis reveals lines of singularities, as shown in Fig. 2. These are obtained consistently for the staggered

magnetization M and the uniform perpendicular susceptibility χ . We note, in particular, that for the critical point x_c is about equal to 1 at $J_2/J_1 = 2.55$, this is consistent with the critical point determined by the dimer expansions.

At the next stage of the analysis, we try to extrapolate the series to the isotropic point ($x = 1$) for those values of the exchange coupling parameters which lie within the Néel-ordered phase at $x = 1$. For this purpose, we first transform the series to a new variable

$$\delta = 1 - (1 - x)^{1/2} \quad (13)$$

to remove the singularity at $x = 1$ predicted by the spin-wave theory. This was first proposed by Huse [17] and was also used in our earlier work on the square lattice case [18]. We then use both the integrated first-order inhomogeneous differential approximants [15] and the naive Padé approximants to extrapolate the series to the isotropic point $\delta = 1$ ($x = 1$). The results for the ground state energy per site E_0/N , the staggered magnetization M , the uniform perpendicular susceptibility $\chi_\perp$, and spin-wave velocity v , and the spin-wave stiffness ρ_s are shown in Figs. 3-5. The results for $J_2 = 0$ presented in the figures are taken from our earlier work on the single-layer square lattice [18,19]. The estimates of the ground state energy per site E_0/N from dimer expansions is also shown in Fig. 3. Note that in the classical spin-wave (large S) level, the uniform perpendicular susceptibility $\chi_\perp$, the spin-wave velocity v , and spin-wave stiffness ρ_s are found to be:

$$\begin{aligned} \chi_\perp^{\text{classical}} &= \frac{1}{8J_1}(1 - y/4) \\ v^{\text{classical}} &= 2\sqrt{2}SJ_1(1 + y/4)^{1/2} \\ \rho_s^{\text{classical}} &= J_1S^2(1 - y^2/16) \end{aligned} \quad (14)$$

so in order to remove the trivial dependent of these quantities on y , Figs. 4-5 have been presented in terms of quantum renormalizations of the classical spin-wave values. We note that M , $\chi_\perp$, and ρ_s first increase for small J_2/J_1 , pass through a maximum at about $y \simeq 0.5$, and then decrease for larger value of J_2/J_1 , and vanishes at about $J_2/J_1 = 2.55$, which is consistent with the critical point determined by the dimer expansion. The reason that in the case of small J_2/J_1 , the interlayer coupling enhances the antiferromagnetic long-range order is the system acquires a weak three dimensionality and quantum fluctuation is suppressed.

The antisymmetric excitation spectra $\Delta(k_x, k_y, \pi)$ for some particular values of y are illustrated in Fig. 6, where we can see that the excitation is gapless at (π, π, π) point. In Fig. 7, we present a plot of energy gaps at $(0, 0, \pi)$ and $(\pi, 0, \pi)$ obtained from the linear spin-wave expansion, the Schwinger boson method [3], the dimer expansions, and the Ising expansions. We can see, from Figs. 6-7, that the spin-wave gap evolves very smoothly from the ordered side to the disordered side of the transition, (the gap at $X = (0, 0, \pi)$ turns out to have a small discontinuity at transition point, which may be due to the estimates from Ising expansions being a little too small). From Fig. 7, we can also see that for Δ_X , the results of leading order spin-wave theory is about 10% smaller than our estimates from Ising expansions. For optical magnon gap Δ_{opt} , the Schwinger boson method give consistent results with those estimated from the Ising expansions over the region $0.2 \lesssim y \lesssim 1$, but as y decreases from 0.2, the optical magnon gap estimated by the Ising expansions becomes larger than that predicted by the spin-wave theory, and also by the Schwinger boson method.

Finally, let us estimate the interlayer exchange constant for $\text{YBa}_2\text{Cu}_3\text{O}_{6.2}$ using the optical magnon gap estimated by Ising expansions: the recent neutron scattering measurements [1,2] have report that the optical magnon gap for $\text{YBa}_2\text{Cu}_3\text{O}_{6.2}$ is $\Delta_{\text{opt}} \simeq 70\text{meV}$, and if we take the canonical value of $J_1 = 120\text{meV}$, we estimate that $J_2/J_1 \simeq 0.07$, whereas J_2/J_1 estimated by the spin-wave theory [1,2] and by the Schwinger boson theory [3] are 0.08 and 0.14, respectively.

IV. HIGH TEMPERATURE SERIES EXPANSIONS

We now turn to the thermodynamic properties of the system at finite temperatures. We have developed high-temperature series expansions for the uniform magnetic susceptibility $\chi(T)$ and the specific heat $C(T)$, for the system with arbitrary J_1/J_2 ,

$$T\chi(T) = \frac{1}{N} \sum_i \sum_j \frac{\text{Tr} S_i^z S_j^z e^{-\beta H}}{\text{Tr} e^{-\beta H}} \quad (15)$$

$$C(T) = \frac{\partial U}{\partial T}$$

where N is the number of sites, and $\beta = J_2/(k_B T)$, and the internal energy U is defined by

$$U = \frac{\text{Tr} H e^{-\beta H}}{\text{Tr} e^{-\beta H}} \quad (16)$$

The series were computed to order β^9 for $\chi(T)$ and to order β^8 for $C(T)$. These series are listed in Table IV. We use the integrated first-order inhomogeneous differential approximants [15] and the naive Padé approximants to extrapolate the series. The resulting estimates for some particular values of J_1/J_2 are shown in Figs. 8-9. For the susceptibility, in order to remove the trivial dependence of $\chi(T)$ on Curie-Weiss temperature, we present in Fig. 8 the $(1/\chi(T) - T_{\text{cw}})/J_1$ as a function of T/J_1 where the Curie-Weiss temperature $T_{\text{cw}} = J_1 + J_2/4$. From this graph, we can see that for $y > y_c$, the susceptibility $\chi(T)$ converge nicely to very low temperature, and vanishes in the low temperature limit, which consistent with a gapped system. For $y \lesssim y_c$, the extrapolation, as judged by the consistency of different approximants, do not converge well in low-temperature region. At the critical ratio $(J_2/J_1)_c$, Monte Carlo simulations [6] have shown that the universal, linear behavior of $\chi(T)$ extends up to very high temperature $T \sim J_1$, unfortunately our series extrapolation is not reliable enough to see this behavior.

V. CONCLUSIONS

In the preceding sections we have presented and analyzed a variety of high-order perturbation series expansions for the two-layer, $S = \frac{1}{2}$, square lattice Heisenberg (and Heisenberg-Ising) model. At $T = 0$, we have added three more terms to existing series for the ground state energy E_0/N , the antiferromagnetic susceptibility χ , and the lowest lying triplet excitation spectrum $\Delta(k_x, k_y)$ from the expansions about the interlayer dimer limits, and also

carried out new perturbation series expansions about the Ising limit up to tenth order for a number of properties. In addition, high-temperature series for the system with arbitrary J_1/J_2 are computed to order β^9 for uniform magnetic susceptibility $\chi(T)$ and to order β^8 for the specific heat $C(T)$.

Based on exponent-biased analyses of the high-order dimer expansions for triplet energy gap and antiferromagnetic susceptibility, we estimate the antiferromagnetic order-disorder transition occurring at $T = 0$ at a critical coupling ratio $(J_2/J_1)_c = 2.537(5)$, which is in good agreement with the quantum Monte Carlo results [6] $(J_2/J_1)_c = 2.51(2)$. The spin-wave velocity at the critical point is estimated to be $v/J_2 = 0.748(2)$, (equivalently, $v/J_1 = 1.898(9)$), which is slightly higher than the QMC results [6] $v/J_1 = 1.69(2)$. The spin-triplet excitation spectra for the system on both ordered and disordered phases are computed, and they seem to be evolve smoothly from the ordered side to the disordered side of the transition. At the limit of $J_2/J_1 \rightarrow 0$, the optical magnon gap obtained by the Ising expansions is larger than $2\sqrt{J_1 J_2}$, instead of less than $2\sqrt{J_1 J_2}$ as predicted by the Schwinger boson method, and this implies the interlayer exchange constant is $J_2 \simeq 0.07J_1$ for $\text{YBa}_2\text{Cu}_3\text{O}_{6.2}$.

ACKNOWLEDGMENTS

This work forms part of a research project supported by a grant from the Australian Research Council. I would like to thank Profs. J. Oitmaa and C.J. Hamer for a number of valuable discussions and suggestions.

REFERENCES

* e-mail address: w.zheng@unsw.edu.au

- [1] D. Reznik, *et. al.*, Phys. Rev. B **53**, 14741(1996).
- [2] S.M. Hayden, G. Aeppli, T.G. Perring, H.A. Mook and F. Dogan, Phys. Rev. B **54**, R6905(1996).
- [3] A.J. Millis and H. Monien, Phys. Rev. B **54**, 16172(1996).
- [4] K. Hida, J. Phys. Soc. Jpn. **61**, 1013(1992).
- [5] M. P. Gelfand, Phys. Rev. B **53**, 11309(1996).
- [6] A.W. Sandvik and D.J. Scalapino, Phys. Rev. Lett. **72**, 2777(1994).
- [7] A.W. Sandvik, A.V. Chubukov, and S. Sachdev, Phys. Rev. B **51**, 16483(1995).
- [8] A.W. Sandvik and D.J. Scalapino, Phys. Rev. B **53**, R526(1996).
- [9] A.J. Millis and H. Monien, Phys. Rev. Lett. **70**, 2810(1993); Phys. Rev. B **50**, 16606(1994).
- [10] A.V. Chubukov and D.K. Morr, Phys. Rev. B **52**, 3521(1995).
- [11] T. Miyazaki, I. Nakamura, and D. Yoshioka, Phys. Rev. B **53**, 12206(1996).
- [12] H.X. He, C.J. Hamer and J. Oitmaa, J. Phys. A **23**, 1775(1990).
- [13] M. P. Gelfand, R.R.P. Singh, and D.A. Huse, J. of Stat. Phys. **59**, 1093(1990).
- [14] M. P. Gelfand, Solid State Commun. **98**, 11(1996).
- [15] A.J. Guttman, in *Phase Transitions and Critical Phenomena*, edited by C. Domb and J.L. Lebowitz (Academic, New York, 1989), Vol. 13.
- [16] T. Matsuda and K. Hida, J. Phys. Soc. Jpn. **59**, 2223(1990); K. Hida, J. Phys. Soc. Jpn. **59**, 2230(1990).
- [17] D.A. Huse, Phys. Rev. B **37**, 2380(1988).
- [18] W.H. Zheng, J. Oitmaa and C.J. Hamer, Phys. Rev. B **43**, 8321(1991).
- [19] C.J. Hamer, W.H. Zheng, and J. Oitmaa, Phys. Rev. B **50**, 6877(1994).

FIGURES

FIG. 1. Plot of the antisymmetric spin-triplet excitation spectrum $\Delta(k_x, k_y)$ along high-symmetry cuts through the Brillouin zone for the system with coupling ratios $J_1/J_2 = 0.1, 0.2, 0.3, 0.3942$ (shown in the figure from the top to the bottom at (π, π) respectively), the lines are the estimates by direct sum to the dimer series, and the points (circles with error bar for the case of $J_1/J_2 = 0.3942$ only) are the estimates of the Padé approximants to the dimer series.

FIG. 2. The phase boundary for the Heisenberg-Ising model without the staggered field ($t = 0$) obtained from analysis of the Ising series for staggered magnetization M and uniform perpendicular susceptibility $\chi_\perp$.

FIG. 3. The rescaled ground-state energy per site E_0/N as function of $J_2/(J_1 + J_2)$. The crosses with dashed line connecting them are the estimates from Ising expansion, and the solid curve is the estimates derived from the dimer expansions. The error bars are much smaller than the symbols.

FIG. 4. The staggered magnetization M and the renormalization factor of the uniform perpendicular susceptibility $Z_\chi = 8\chi_\perp J_1/(1 - y/4)$ (at $T = 0$) *versus* J_2/J_1 as estimated by Ising expansions.

FIG. 5. The renormalization factor of the spin-wave velocity $Z_c = v/[J_1(2 + y/2)^{1/2}]$ and the spin-wave stiffness $Z_\rho = 4\rho_s/[J_1(1 - y^2/16)]$ *versus* J_2/J_1 as estimated by Ising expansions and dimer expansions (for Z_c at the critical ratio $(J_2/J_1)_c = 2.537$ only).

FIG. 6. Plot of the antisymmetric spin-triplet excitation spectrum $\Delta(k_x, k_y, \pi)$ (derived from the Ising expansions) along high-symmetry cuts through the Brillouin zone for the system with coupling ratios $J_2/J_1 = 0.5, 1, 1.5, 2.5$ (shown in the figure from the top to the bottom at $(\pi, 0)$ respectively)

FIG. 7. The rescaled energy gap at $(0, 0, \pi)$ (the optical magnon gap) and at $X = (\pi, 0, \pi)$ as function of $J_2/(J_1 + J_2)$. The solid curves at large $J_2/(J_1 + J_2)$ are the extrapolation derived from the dimer expansions, the crosses with error bars are the estimates of Ising expansions, the long dashed lines at small $J_2/(J_1 + J_2)$ are the results of the linear spin-wave theory, the open circles connected by a short dashed line are the results of Eq.(12) (where the ρ_s is taken from the results of Fig. 5), and the vertical dotted line indicates the position of the transition.

FIG. 8. Susceptibility $(1/\chi(T) - T_{cw})/J_1$ as a function of temperature for the system with coupling ratios $J_2/J_1 = 5, 10/3, 2.537, 5/3, 1$, several different integrated differential approximants to the high temperature series are shown for each J_2/J_1 .

FIG. 9. The specific heat $C(T)$ as a function of temperature for the system with coupling ratios $J_1/J_2 = 0.1, 0.3$, several different integrated differential approximants to the high temperature series are shown for each J_1/J_2

TABLES

TABLE I. Series coefficients for dimer expansions of the ground-state energy per site $E_0/(NJ_2)$, the antiferromagnet susceptibility χ , the gap m/J_2 , and the critical spin-wave velocity (as described in the text). Coefficients of $(1/y)^n$ up to order $n = 11$ are listed.

n	$E_0/(NJ_2)$	$J_2\chi$	m/J_2	$2CD/J_2^2$
0	$-3.750000000 \times 10^{-1}$	1.000000000	1.000000000	0.000000000
1	0.000000000	4.000000000	-2.000000000	1.000000000
2	$-3.750000000 \times 10^{-1}$	1.300000000×10^1	0.000000000	0.000000000
3	$-1.875000000 \times 10^{-1}$	3.950000000×10^1	-1.500000000	2.375000000
4	$-1.171875000 \times 10^{-1}$	1.128125000×10^2	-1.500000000	1.437500000
5	$1.289062500 \times 10^{-1}$	3.131076389×10^2	$-5.468750000 \times 10^{-1}$	-1.320312500
6	$2.373046875 \times 10^{-1}$	8.535167824×10^2	$1.191406250 \times 10^{-1}$	$-7.851562500 \times 10^{-1}$
7	$2.162475586 \times 10^{-1}$	2.304592044×10^3	-4.202636719	-2.716064453
8	$-4.346008301 \times 10^{-1}$	6.169223926×10^3	-1.060513306×10^1	3.307617188
9	$-9.492845535 \times 10^{-1}$	1.640000304×10^4	-2.693954849×10^1	1.409708977×10^1
10	-1.231647968	4.332290008×10^4	-3.315443579×10^1	3.259252389
11	$6.411397507 \times 10^{-1}$	1.139206022×10^5	-8.044747781×10^1	2.037096158×10^1

TABLE II. Series coefficients for Ising expansions of the ground-state energy per site $E_0/(NJ_1)$, the staggered magnetization M , uniform perpendicular susceptibility $\chi_\perp$, the energy gap $m = \Delta(0,0,0) = \Delta(\pi,\pi,\pi)$, and the critical spin-wave velocity $2CD$ (as described in the text) for $J_2/J_1 = 0.5, 1, 1.5, 2.5$ and $t = 0$, Nonzero coefficients x^n up to order $n = 10$ are listed.

n	$J_2/J_1 = 0.5$	$J_2/J_1 = 1$	$J_2/J_1 = 1.5$	$J_2/J_1 = 2.5$
$E_0/(J_1 N)$				
0	$-5.62500000 \times 10^{-1}$	$-6.25000000 \times 10^{-1}$	$-6.87500000 \times 10^{-1}$	$-8.12500000 \times 10^{-1}$
2	$-1.50669643 \times 10^{-1}$	$-1.56250000 \times 10^{-1}$	$-1.81423611 \times 10^{-1}$	$-2.86221591 \times 10^{-1}$
4	$1.06782048 \times 10^{-3}$	$2.32514881 \times 10^{-5}$	$-1.82797961 \times 10^{-3}$	$-1.01025331 \times 10^{-2}$
6	$-1.55432939 \times 10^{-3}$	$-2.51031531 \times 10^{-3}$	$-3.56180060 \times 10^{-3}$	$-5.93087945 \times 10^{-3}$
8	$-6.14963212 \times 10^{-4}$	$-8.46200313 \times 10^{-4}$	$-1.01380621 \times 10^{-3}$	$-1.82417072 \times 10^{-3}$
10	$-3.42105908 \times 10^{-4}$	$-5.02769527 \times 10^{-4}$	$-6.24621980 \times 10^{-4}$	$-1.03444977 \times 10^{-3}$
$J_1 \chi_\perp$				
0	$2.22222222 \times 10^{-1}$	$2.00000000 \times 10^{-1}$	$1.81818182 \times 10^{-1}$	$1.53846154 \times 10^{-1}$
1	$-2.81746032 \times 10^{-1}$	$-2.50000000 \times 10^{-1}$	$-2.29797980 \times 10^{-1}$	$-2.08041958 \times 10^{-1}$
2	$2.92713075 \times 10^{-1}$	$2.59090909 \times 10^{-1}$	$2.37859757 \times 10^{-1}$	$2.10928368 \times 10^{-1}$
3	$-3.04452465 \times 10^{-1}$	$-2.68827110 \times 10^{-1}$	$-2.49554660 \times 10^{-1}$	$-2.34325217 \times 10^{-1}$
4	$3.05483508 \times 10^{-1}$	$2.69863988 \times 10^{-1}$	$2.49828011 \times 10^{-1}$	$2.29571097 \times 10^{-1}$
5	$-3.09716015 \times 10^{-1}$	$-2.74883091 \times 10^{-1}$	$-2.56566684 \times 10^{-1}$	$-2.43161906 \times 10^{-1}$
6	$3.10814223 \times 10^{-1}$	$2.76075768 \times 10^{-1}$	$2.57806919 \times 10^{-1}$	$2.44310355 \times 10^{-1}$
7	$-3.13649358 \times 10^{-1}$	$-2.79129287 \times 10^{-1}$	$-2.61531350 \times 10^{-1}$	$-2.53169737 \times 10^{-1}$
8	$3.14366380 \times 10^{-1}$	$2.79764322 \times 10^{-1}$	$2.62062184 \times 10^{-1}$	$2.52501211 \times 10^{-1}$
9	$-3.16219779 \times 10^{-1}$	$-2.81901242 \times 10^{-1}$	$-2.64657763 \times 10^{-1}$	$-2.58530073 \times 10^{-1}$
M				
0	$5.00000000 \times 10^{-1}$	$5.00000000 \times 10^{-1}$	$5.00000000 \times 10^{-1}$	$5.00000000 \times 10^{-1}$
2	$-8.55389031 \times 10^{-2}$	$-7.81250000 \times 10^{-2}$	$-8.45389660 \times 10^{-2}$	$-1.30714101 \times 10^{-1}$
4	$-8.46292613 \times 10^{-3}$	$-8.53938581 \times 10^{-3}$	$-1.34925210 \times 10^{-2}$	$-4.35583219 \times 10^{-2}$
6	$-5.85842608 \times 10^{-3}$	$-8.07527219 \times 10^{-3}$	$-1.15477220 \times 10^{-2}$	$-2.85436734 \times 10^{-2}$
8	$-3.93110760 \times 10^{-3}$	$-4.86105744 \times 10^{-3}$	$-6.30805266 \times 10^{-3}$	$-1.93128872 \times 10^{-2}$
10	$-2.73750635 \times 10^{-3}$	$-3.51123409 \times 10^{-3}$	$-4.48806496 \times 10^{-3}$	$-1.37095158 \times 10^{-2}$
m/J_1				
0	2.25000000	2.50000000	2.75000000	3.25000000
2	-1.64330357	-1.77083333	-1.98338294	-2.55445076
4	$1.20240301 \times 10^{-1}$	$5.95393105 \times 10^{-2}$	$9.76911256 \times 10^{-2}$	$3.91257865 \times 10^{-1}$
6	$-2.91150072 \times 10^{-1}$	$-2.85968590 \times 10^{-1}$	$-3.69547989 \times 10^{-1}$	$-9.18026495 \times 10^{-1}$
8	$8.98232628 \times 10^{-2}$	$6.28691338 \times 10^{-2}$	$1.56753567 \times 10^{-1}$	1.10864903
10	$-1.90221034 \times 10^{-1}$	$-1.73683175 \times 10^{-1}$	$-3.02886458 \times 10^{-1}$	-2.00750062
$2CD/J_1^2$				
2	3.97500000	4.16666667	4.51785714	5.59722222
4	-1.62276149	-1.51750579	-1.78959346	-3.20894532
6	1.30765116	1.08647250	1.39046692	3.71475815
8	-1.40693583	-1.13102684	-1.70310223	-7.26139043
10	1.60579536	1.23771527	2.19444652	1.47488833×10^1

TABLE III. Series coefficients for the Ising expansions of the triplet spin-wave excitation spectrum $\Delta(k_x, k_y, k_z) = J_1 \sum_{k,n,m,p} a_{k,n,m,p} x^k \cos(pk_z) [\cos(mk_x) \cos(nk_y) + \cos(nk_x) \cos(mk_y)]/2$. Nonzero coefficients $a_{k,n,m,p}$ up to order $k = 10$ for the case of $J_2 = J_1$ are listed.

(k,n,m,p)	$a_{k,n,m,p}$	(k,n,m,p)	$a_{k,n,m,p}$	(k,n,m,p)	$a_{k,n,m,p}$	(k,n,m,p)	$a_{k,n,m,p}$
(0,0,0,0)	2.50000000	(6,1,1,0)	$-5.01421733 \times 10^{-2}$	(6,3,3,0)	$-3.50360695 \times 10^{-3}$	(10,7,0,1)	$-1.95483982 \times 10^{-4}$
(2,0,0,0)	$-1.04166667 \times 10^{-1}$	(8,1,1,0)	$3.13641370 \times 10^{-2}$	(8,3,3,0)	$-2.76765928 \times 10^{-3}$	(10,9,0,1)	$-3.18431828 \times 10^{-6}$
(4,0,0,0)	$8.82430142 \times 10^{-2}$	(10,1,1,0)	$-2.49468947 \times 10^{-2}$	(10,3,3,0)	$-2.34255921 \times 10^{-3}$	(4,2,1,1)	$-1.18055556 \times 10^{-1}$
(6,0,0,0)	$-2.63720521 \times 10^{-2}$	(4,3,1,0)	$-3.93518519 \times 10^{-2}$	(8,5,3,0)	$-7.65220675 \times 10^{-4}$	(6,2,1,1)	$-2.64155861 \times 10^{-2}$
(8,0,0,0)	$1.81429057 \times 10^{-2}$	(6,3,1,0)	$-1.87035451 \times 10^{-2}$	(10,5,3,0)	$-1.48448940 \times 10^{-3}$	(8,2,1,1)	$5.61540297 \times 10^{-3}$
(10,0,0,0)	$-9.88646062 \times 10^{-3}$	(8,3,1,0)	$-3.19610735 \times 10^{-3}$	(10,7,3,0)	$-7.64236388 \times 10^{-5}$	(10,2,1,1)	$-2.48059104 \times 10^{-2}$
(2,2,0,0)	$-3.33333333 \times 10^{-1}$	(10,3,1,0)	$-1.18183470 \times 10^{-2}$	(8,4,4,0)	$-4.78262922 \times 10^{-4}$	(6,4,1,1)	$-1.05108208 \times 10^{-2}$
(4,2,0,0)	$1.52199074 \times 10^{-2}$	(6,5,1,0)	$-2.10216417 \times 10^{-3}$	(10,4,4,0)	$-8.91369509 \times 10^{-4}$	(8,4,1,1)	$-5.39907287 \times 10^{-3}$
(6,2,0,0)	$-3.34161849 \times 10^{-2}$	(8,5,1,0)	$-2.49774309 \times 10^{-3}$	(10,6,4,0)	$-1.33741368 \times 10^{-4}$	(10,4,1,1)	$-5.30710289 \times 10^{-3}$
(8,2,0,0)	$1.25625535 \times 10^{-2}$	(10,5,1,0)	$-2.84123845 \times 10^{-3}$	(10,5,5,0)	$-8.02448208 \times 10^{-5}$	(8,6,1,1)	$-7.65220675 \times 10^{-4}$
(10,2,0,0)	$-1.75733942 \times 10^{-2}$	(8,7,1,0)	$-1.09317239 \times 10^{-4}$	(2,1,0,1)	$-6.66666667 \times 10^{-1}$	(10,6,1,1)	$-1.16719696 \times 10^{-3}$
(4,4,0,0)	$-4.91898148 \times 10^{-3}$	(10,7,1,0)	$-3.13087047 \times 10^{-4}$	(4,1,0,1)	$6.74768519 \times 10^{-2}$	(10,8,1,1)	$-5.73177291 \times 10^{-5}$
(6,4,0,0)	$-4.91156561 \times 10^{-3}$	(10,9,1,0)	$-6.36863657 \times 10^{-6}$	(6,1,0,1)	$-5.84006701 \times 10^{-2}$	(6,3,2,1)	$-2.10216417 \times 10^{-2}$
(8,4,0,0)	$-2.30378946 \times 10^{-3}$	(4,2,2,0)	$-2.95138889 \times 10^{-2}$	(8,1,0,1)	$3.34318983 \times 10^{-2}$	(8,3,2,1)	$-5.04717657 \times 10^{-3}$
(10,4,0,0)	$-3.17090252 \times 10^{-3}$	(6,2,2,0)	$-1.09825781 \times 10^{-2}$	(10,1,0,1)	$-3.12876108 \times 10^{-2}$	(10,3,2,1)	$-7.50314475 \times 10^{-3}$
(6,6,0,0)	$-1.75180347 \times 10^{-4}$	(8,2,2,0)	$-7.99539319 \times 10^{-4}$	(4,3,0,1)	$-1.96759259 \times 10^{-2}$	(8,5,2,1)	$-2.29566203 \times 10^{-3}$
(8,6,0,0)	$-3.97833036 \times 10^{-4}$	(10,2,2,0)	$-7.65794740 \times 10^{-3}$	(6,3,0,1)	$-1.30043279 \times 10^{-2}$	(10,5,2,1)	$-2.08196519 \times 10^{-3}$
(10,6,0,0)	$-6.37350643 \times 10^{-4}$	(6,4,2,0)	$-5.25541042 \times 10^{-3}$	(8,3,0,1)	$-9.25242372 \times 10^{-4}$	(10,7,2,1)	$-2.29270916 \times 10^{-4}$
(8,8,0,0)	$-6.83232746 \times 10^{-6}$	(8,4,2,0)	$-4.57905354 \times 10^{-3}$	(10,3,0,1)	$-6.65895455 \times 10^{-3}$	(8,4,3,1)	$-3.82610338 \times 10^{-3}$
(10,8,0,0)	$-3.37994352 \times 10^{-5}$	(10,4,2,0)	$-4.19352456 \times 10^{-3}$	(6,5,0,1)	$-1.05108208 \times 10^{-3}$	(10,4,3,1)	$-2.63082695 \times 10^{-3}$
(10,10,0,0)	$-3.18431828 \times 10^{-7}$	(8,6,2,0)	$-3.82610338 \times 10^{-4}$	(8,5,0,1)	$-1.65065868 \times 10^{-3}$	(10,6,3,1)	$-5.34965472 \times 10^{-4}$
(2,1,1,0)	$-6.66666667 \times 10^{-1}$	(10,6,2,0)	$-8.44976702 \times 10^{-4}$	(10,5,0,1)	$-1.45569445 \times 10^{-3}$	(10,5,4,1)	$-8.02448208 \times 10^{-4}$
(4,1,1,0)	$1.00115741 \times 10^{-1}$	(10,8,2,0)	$-2.86588646 \times 10^{-5}$	(8,7,0,1)	$-5.46586197 \times 10^{-5}$		

TABLE IV. Series coefficients for the high-temperature expansions of the uniform susceptibility $\chi(T) = T^{-1} \sum_{m,n} a_{m,n} (J_1/J_2)^m \beta^n$ and the specific heat $C(T) = \beta^2 \sum_{m,n} a_{m,n} (J_1/J_2)^m \beta^n$. Nonzero coefficients $a_{m,n}$ up to order $n = 9$ for χ or order $n = 8$ for C are listed.

(m,n)	$a_{m,n}$	(m,n)	$a_{m,n}$	(m,n)	$a_{m,n}$	(m,n)	$a_{m,n}$
$\chi(T)$							
(0,0)	$2.500000000 \times 10^{-1}$	(4,4)	$1.692708333 \times 10^{-2}$	(0,7)	$-3.986661396 \times 10^{-5}$	(6,8)	$1.061827644 \times 10^{-2}$
(0,1)	$-6.250000000 \times 10^{-2}$	(0,5)	$2.115885417 \times 10^{-4}$	(1,7)	$5.113389757 \times 10^{-4}$	(7,8)	$4.403056796 \times 10^{-3}$
(1,1)	$-2.500000000 \times 10^{-1}$	(1,5)	$-3.580729167 \times 10^{-3}$	(2,7)	$-2.441406250 \times 10^{-3}$	(8,8)	$3.978426494 \times 10^{-4}$
(0,2)	$-1.562500000 \times 10^{-2}$	(2,5)	$1.822916667 \times 10^{-2}$	(3,7)	$4.155815972 \times 10^{-3}$	(0,9)	$3.899804709 \times 10^{-6}$
(1,2)	$1.250000000 \times 10^{-1}$	(3,5)	$-2.864583333 \times 10^{-2}$	(4,7)	$6.287977431 \times 10^{-3}$	(1,9)	$-4.881601485 \times 10^{-5}$
(2,2)	$1.250000000 \times 10^{-1}$	(4,5)	$-4.492187500 \times 10^{-2}$	(5,7)	$-1.420898438 \times 10^{-2}$	(2,9)	$2.027723524 \times 10^{-4}$
(0,3)	$1.302083333 \times 10^{-3}$	(5,5)	$-9.244791667 \times 10^{-3}$	(6,7)	$-1.158582899 \times 10^{-2}$	(3,9)	$-1.131895358 \times 10^{-4}$
(1,3)	$1.562500000 \times 10^{-2}$	(0,6)	$-1.044379340 \times 10^{-4}$	(7,7)	$6.285652282 \times 10^{-4}$	(4,9)	$-1.836698017 \times 10^{-3}$
(2,3)	$-1.250000000 \times 10^{-1}$	(1,6)	$5.533854167 \times 10^{-4}$	(0,8)	$1.513768756 \times 10^{-6}$	(5,9)	$3.601243761 \times 10^{-3}$
(3,3)	$-4.166666667 \times 10^{-2}$	(2,6)	$1.399739583 \times 10^{-3}$	(1,8)	$3.700861855 \times 10^{-5}$	(6,9)	$3.023339327 \times 10^{-4}$
(0,4)	$1.627604167 \times 10^{-3}$	(3,6)	$-1.527777778 \times 10^{-2}$	(2,8)	$-6.515260727 \times 10^{-4}$	(7,9)	$-7.574656653 \times 10^{-3}$
(1,4)	$-1.041666667 \times 10^{-2}$	(4,6)	$2.438151042 \times 10^{-2}$	(3,8)	$3.424169147 \times 10^{-3}$	(8,9)	$-1.439339774 \times 10^{-3}$
(2,4)	$7.812500000 \times 10^{-3}$	(5,6)	$2.350260417 \times 10^{-2}$	(4,8)	$-5.407569522 \times 10^{-3}$	(9,9)	$-5.211347415 \times 10^{-4}$
(3,4)	$8.333333333 \times 10^{-2}$	(6,6)	$1.991102431 \times 10^{-3}$	(5,8)	$-1.251705109 \times 10^{-3}$		
$C(T)$							
(0,0)	$9.375000000 \times 10^{-2}$	(2,4)	$-1.025390625 \times 10^{-2}$	(3,6)	$3.450520833 \times 10^{-3}$	(9,7)	$-6.530006045 \times 10^{-2}$
(2,0)	$3.750000000 \times 10^{-1}$	(3,4)	$-1.367187500 \times 10^{-2}$	(4,6)	$-4.157307943 \times 10^{-2}$	(0,8)	$-5.264736357 \times 10^{-5}$
(0,1)	$4.687500000 \times 10^{-2}$	(4,4)	$1.464843750 \times 10^{-1}$	(5,6)	$2.871093750 \times 10^{-2}$	(2,8)	$-5.870092483 \times 10^{-4}$
(3,1)	$1.875000000 \times 10^{-1}$	(6,4)	$3.173828125 \times 10^{-2}$	(6,6)	$-8.115234375 \times 10^{-2}$	(3,8)	$-4.486810593 \times 10^{-4}$
(0,2)	$-5.859375000 \times 10^{-3}$	(0,5)	$9.399414063 \times 10^{-4}$	(8,6)	$1.951700846 \times 10^{-2}$	(4,8)	$3.243491763 \times 10^{-3}$
(2,2)	$-4.687500000 \times 10^{-2}$	(2,5)	$4.785156250 \times 10^{-3}$	(0,7)	$-1.816522507 \times 10^{-5}$	(5,8)	$-1.445297968 \times 10^{-2}$
(4,2)	$-1.640625000 \times 10^{-1}$	(3,5)	$3.759765625 \times 10^{-3}$	(2,7)	$7.376534598 \times 10^{-4}$	(6,8)	$2.286320641 \times 10^{-2}$
(0,3)	$-9.765625000 \times 10^{-3}$	(4,5)	$4.990234375 \times 10^{-2}$	(3,7)	$2.947126116 \times 10^{-4}$	(7,8)	$-3.248014904 \times 10^{-2}$
(2,3)	$-3.906250000 \times 10^{-2}$	(5,5)	$9.365234375 \times 10^{-2}$	(4,7)	$-2.781633650 \times 10^{-2}$	(8,8)	$3.321502322 \times 10^{-2}$
(3,3)	$-3.906250000 \times 10^{-2}$	(7,5)	$1.268066406 \times 10^{-1}$	(5,7)	$-2.221854074 \times 10^{-2}$	(10,8)	$-2.914089021 \times 10^{-2}$
(5,3)	$-1.953125000 \times 10^{-1}$	(0,6)	$4.185994466 \times 10^{-4}$	(6,7)	$-2.977120536 \times 10^{-2}$		
(0,4)	$-1.586914062 \times 10^{-3}$	(2,6)	$4.451497396 \times 10^{-3}$	(7,7)	$-1.089634487 \times 10^{-1}$		

















